

Dual-antiplatelet treatment beyond 1 year after drug-eluting stent implantation (ARCTIC-Interruption): a randomised trial



Jean-Philippe Collet, Johanne Silvain, Olivier Barthélémy, Grégoire Rangé, Guillaume Cayla, Eric Van Belle, Thomas Cuisset, Simon Elhadad, François Schiele, Nicolas Lhoest, Patrick Ohlmann, Didier Carrié, Hélène Rousseau, Pierre Aubry, Jacques Monségu, Pierre Sabouret, Stephen A O'Connor, Jérémie Abtan, Mathieu Kerneis, Christophe Saint-Etienne, Farzin Beygui, Eric Vicaut, Gilles Montalescot, for the ARCTIC investigators*

Summary

Background Optimum duration of dual antiplatelet treatment (DAPT) after coronary stenting remains uncertain, with an unknown efficacy to safety ratio of extended treatment leading to discrepancies between international guidelines and clinical practice. We assessed whether DAPT continuation beyond 1 year after coronary stenting is beneficial.

Methods This analysis was a planned extension of the previously published ARCTIC-Monitoring trial, in which we randomly allocated 2440 patients to a strategy of platelet function testing with antiplatelet treatment adjustment or a conventional strategy after coronary stenting with drug-eluting stent (DES). We recruited patients (aged 18 years or older) scheduled for planned DES implantation at 38 centres in France. After 1 year of follow-up, patients without contraindication to interruption of DAPT were eligible for a second randomisation to this second phase of the study (ARCTIC-Interruption). Using a computer-generated randomisation sequence (1:1; stratified by centre), we allocated patients to a strategy of interruption of DAPT where the thienopyridine was interrupted and single aspirin antiplatelet treatment was maintained (interruption group) or a strategy of DAPT continuation for 6–18 months (continuation group). The primary endpoint was the composite of death, myocardial infarction, stent thrombosis, stroke, or urgent revascularisation, analysed by intention to treat. This trial is registered with ClinicalTrials.gov, number NCT00827411.

Findings Between Jan 4, 2011, and March 3, 2012, 1259 eligible patients were randomly allocated to treatment in ARCTIC-Interruption: 624 to the interruption group and 635 to the continuation group. After a median follow-up of 17 months (IQR 15–18), the primary endpoint occurred in 27 (4%) patients in the interruption group and 24 (4%) patients in the continuation group (hazard ratio [HR] 1.17 [95% CI 0.68–2.03]; $p=0.58$). STEEPLE major bleeding events occurred more often in the continuation group (seven [1%] patients) compared with the interruption group (one [$<0.5\%$] patient; HR 0.15 [0.02–1.20]; $p=0.073$). Major or minor bleedings were also more common in the continuation group compared with the interruption group (12 [2%] patients vs three [1%] patients; HR 0.26 [0.07–0.91]; $p=0.04$).

Interpretation Our finding suggests no apparent benefit but instead harm with extension of DAPT beyond 1 year after stenting with DES when no event has occurred within the first year after stenting. No conclusion can be drawn for high-risk patients who could not be randomised. The consistency between findings from all trials of such interruption suggests the need for a reappraisal of guidelines for DAPT after coronary stenting towards shorter duration of treatment.

Funding Allies in Cardiovascular Trials Initiatives and Organized Networks (ACTION Study Group), Fondation de France, Sanofi-Aventis, Cordis, Medtronic, Boston Scientific, Fondation SGAM.

Introduction

The recommended duration of dual antiplatelet treatment (DAPT) for elective coronary implantation of a bare-metal stent is at least 1 month on the basis of observational data.¹ Concerns about a potential increased risk of stent thrombosis with drug-eluting stents (DESs)^{2–4} led to extension of DAPT from 2 months to 6–12 months.^{5,6} However, new analyses of data from randomised trials⁷ and data from studies of the newer generations of DESs have not substantiated any increased risk, but have instead shown a lower risk of stent thrombosis in higher risk clinical situations such as in ST-elevation myocardial infarction.^{8–10}

Findings from several randomised studies and large observational registries have challenged the recom-

mendation of 6–12 months of DAPT treatment after implantation of a DES in stable patients, although duration in the control groups of these studies was variable and not always reflecting the 12-month recommendation of DAPT duration after DES implantation.^{11–20}

The ARCTIC (Assessment by a double Randomisation of a Conventional antiplatelet strategy versus a monitoring-guided strategy for drug-eluting stent implantation and, of Treatment Interruption versus Continuation 1 year after stenting)-Interruption trial is the second phase of the previously published ARCTIC-Monitoring study,^{21,22} in which we randomly allocated 2440 patients to either platelet function testing with antiplatelet treatment adjustment or conventional antiplatelet treatment after coronary stenting with a DES.

Published Online

July 16, 2014

[http://dx.doi.org/10.1016/S0140-6736\(14\)60612-7](http://dx.doi.org/10.1016/S0140-6736(14)60612-7)

[http://dx.doi.org/10.1016/S0140-6736\(14\)60612-7](http://dx.doi.org/10.1016/S0140-6736(14)60612-7)

See Online/Comment

[http://dx.doi.org/10.1016/S0140-6736\(14\)60612-7](http://dx.doi.org/10.1016/S0140-6736(14)60612-7)

[http://dx.doi.org/10.1016/S0140-6736\(14\)60612-7](http://dx.doi.org/10.1016/S0140-6736(14)60612-7)

*Investigators listed in appendix

ACTION Study Group,

Institut de Cardiologie Hôpital

Pitié-Salpêtrière (APHP),

Université Paris 6, INSERM,

Paris, France (Prof J-P Collet MD,

J Silvain MD, S A O'Connor MD,

O Barthélémy MD,

P Sabouret MD, J Abtan MD,

M Kerneis MD,

Prof G Montalescot MD);

Les Hôpitaux de Chartres,

Le Coudray, France

(G Rangé MD); ACTION Study

Group, Cardiologie, CHU

Carémeau, Nîmes, France

(Prof G Cayla MD); CHRU de

Lille, Lille, France

(H Rousseau MSc,

Prof E Van Belle MD);

Département de Cardiologie,

CHU La Timone, Marseille,

France (Prof T Cuisset MD);

Cardiologie, CH de Lagny-

Marne-la-Vallée, Lagny-sur-

Marne, France (S Elhadad MD);

CHU Jean Minjoz, Besançon,

France (Prof F Schiele MD);

GH du Centre Alsace, France

(N Lhoest MD); CHR

Strasbourg, France

(Prof P Ohlmann MD); CHU

Rangueil, Toulouse, France

(Prof D Carrié MD); ACTION

Study Group, Unité de

Recherche Clinique-Hôpital

Lariboisière (APHP), and

Université Denis Diderot, Paris,

France (H Rousseau,

Prof E Vicaut MD); Centre

Hospitalier Bichat (APHP),

Paris, France (P Aubry MD);

Institut Mutualiste

Montsouris, Paris, France

(Prof J Monségu MD); CHU

Trousseau, Tours, France

(C Saint-Etienne MD); and

ACTION study Group, CHU

Caen, France

(Prof F Beygui, MD)

Correspondence to: Prof Gilles Montalescot, Institut de Cardiologie, Bureau 2-236, Centre Hospitalier Universitaire Pitié-Salpêtrière, 47 Boulevard de l'Hôpital, 75013 Paris, France gilles.montalescot@psl.aphp.fr

After 1 year of follow-up, we invited patients without major events and who were treated with clopidogrel or prasugrel in addition to aspirin to enrol in a second randomisation to either interruption of DAPT as recommended by international guidelines or continuation of DAPT for a further 6–18 months.^{6,23} All patients participated in the first phase of the ARCTIC study, which served as a screening log book for the second randomisation 1 year later to assess this second hypothesis about DAPT duration.

Methods

Study design and population

ARCTIC was a multicentre, prospective open-label study with parallel trial arms and double randomisation. Between Jan 26, 2009, and Jan 5, 2011, we recruited patients (aged ≥18 years) scheduled for planned DES implantation at 38 centres in France.²¹ We excluded patients with primary percutaneous coronary intervention for ST-elevation myocardial infarction, planned use of glycoprotein IIb/IIa inhibitors, chronic anticoagulation treatment, or bleeding diathesis. Results from the first phase (ARCTIC-Monitoring) have been published previously.²² In the first phase, we randomly allocated eligible patients to either platelet function

assessment with adjustment of antiplatelet drugs and doses in patients with inadequate platelet inhibitory response (monitoring group) or treatment without platelet function assessment (conventional group)—we saw no difference between the two groups after 1 year of follow-up.²²

At the end of this follow-up period, and in the absence of contraindication to interruption of DAPT, patients were re-randomised from Jan 4, 2010, to March 3, 2012, to interrupt or to continue thienopyridine according to the initial signed consent form.

In the interruption group, the thienopyridine (clopidogrel or prasugrel) was interrupted and aspirin was maintained as single antiplatelet treatment. The final clinical assessment was done 6 months after the second randomisation of the last patient. In the continuation group, clopidogrel or prasugrel were maintained until the end of the study, with a minimum additional follow-up of 6 months for the last randomised patient, whereas all preceding patients had an accrual follow-up (last follow-up in Jan 26, 2013). In the continuation group, DAPT was maintained at the same dose regimen as during the first year of follow-up.

Exclusion criteria for the second randomisation were the occurrence of any ischaemic event of the primary




Figure 1: Trial profile

*Median time to withdrawal 318 days (IQR 189–392). †Median time to withdrawals 315 days (189–380).

	Continuation (n=635)	Interruption (n=624)
Age in years	64 (57-73)	64 (57-73)
Older than 75 years	117 (18%)	103 (17%)
Women	127 (20%)	121 (19%)
Body weight index (kg/m ²)	27 (25-29)	27 (25-30)
Diabetes	198 (31%)	222 (36%)
Dyslipidaemia	428 (67%)	426 (68%)
Hypertension	376 (59%)	388 (62%)
Current smoker	147 (23%)	152 (24%)
High on-treatment platelet reactivity*	100 (19%)	81 (16%)
Prior stroke	28 (4%)	38 (6%)
Prior heart failure	20 (3%)	23 (4%)
Prior myocardial infarction	197 (31%)	186 (30%)
Prior percutaneous coronary intervention	273 (43%)	249 (40%)
Prior coronary artery bypass graft	47 (7%)	35 (6%)
ACE inhibitors	334 (53%)	322 (52%)
β blockers	389 (61%)	364 (58%)
Statin	428 (67%)	416 (67%)
Proton-pump inhibitors	208 (33%)	183 (29%)
Calcium-channel inhibitor	150 (24%)	116 (19%)
Drug-eluting stent implanted	623 (98%)	618 (99%)
First-generation stent†	270 (43%)	250 (40%)
Second-generation stent	392 (62%)	396 (64%)
Stented vessel		
Left main	18 (3%)	23 (4%)
Left anterior descending	342 (54%)	325 (52%)
Circumflex	209 (33%)	181 (29%)
Right coronary artery	191 (30%)	222 (36%)
Coronary artery bypass graft	7 (1%)	6 (1%)

Data are mean (IQR) or n (%). ACE=angiotensin-converting-enzyme. *Platelet reaction unit of 235 or more using the VerifyNow-P2Y₁₂ assay. †Paclitaxel and sirolimus drug-eluting stent.

Table 1: Demographic, clinical, and procedural baseline characteristics

endpoint or any event of the primary safety endpoint during the first year of follow-up after the randomisation in the first phase of the study, any new revascularisation needing DAPT extension, any contraindication to aspirin continuation such as bleeding gastrointestinal ulcer, or aspirin resistance. Physician's choice not to randomise after 1 year was another exclusion criterion and was recorded in the case report form. We also recorded patients' last platelet function test (VerifyNowP2Y₁₂ assay, Accumetrics, CA, USA) done on the maintenance dose of thienopyridine as a baseline characteristic.

Randomisation and masking

The randomisation list was generated by an independent statistician according to the procedures of the Direction de la Recherche et du Développement de l'Assistance Publique des Hôpitaux de Paris with SAS software (version 9.2) and was stratified by centre. Group allocation was done using an interactive voice response system with a 1:1 ratio.

	Continuation (n=635)	Interruption (n=624)	p value
Treatment at randomisation			
Clopidogrel maintenance dose	569 (90%)	562 (90%)	0.79
Clopidogrel maintenance dose of 75 mg	507 (89%)	482 (86%)	0.039
Clopidogrel maintenance dose of 150 mg	56 (10%)	78 (14%)	0.039
Clopidogrel maintenance dose greater than 150 mg	6 (1%)	2 (<0.5%)	0.039
Prasugrel maintenance dose of 10 mg	54 (9%)	53 (9%)	0.99
Aspirin maintenance dose	628 (99%)	622 (100%)	0.18
Treatment at last follow-up visit			
Clopidogrel maintenance dose	454 (72%)	95 (15%)	<0.0001
Aspirin maintenance dose	597 (94%)	605 (97%)	0.0121
Prasugrel maintenance dose of 10 mg	36 (6%)	14 (2%)	0.0019

Table 2: Use of antiplatelet drugs during trial

	Continuation (n=635)	Interruption (n=624)	Hazard ratio (95% CI)	p value
Any death, myocardial infarction, stent thrombosis, stroke or TIA, urgent revascularisation (primary endpoint)	24 (4%)	27 (4%)	1.17 (0.68-2.03)	0.58
Stent thrombosis (revascularised or not) or any urgent revascularisation (main secondary endpoint)	8 (1%)	10 (2%)	1.30 (0.51-3.30)	0.58
Any death, recurrent acute coronary syndrome, stroke or TIA	21 (3%)	24 (4%)	1.19 (0.66-2.13)	0.56
Death or resuscitated cardiac arrest	7 (1%)	9 (1%)	1.32 (0.49-3.55)	0.58
Death or myocardial infarction	14 (2%)	17 (3%)	1.26 (0.62-2.55)	0.52
Any death, myocardial infarction, stent thrombosis (revascularised or not), stroke or TIA, urgent revascularisation, TIMI major bleed (net clinical benefit)	30 (5%)	28 (5%)	0.97 (0.58-1.62)	0.90
Any death	7 (1%)	9 (1%)	1.32 (0.49-3.55)	0.58
Myocardial infarction	9 (1%)	9 (1%)	1.04 (0.41-2.62)	0.94
Stent thrombosis*	0 (0%)	3 (1%)		
Acute coronary syndrome	11 (2%)	13 (2%)	1.23 (0.55-2.74)	0.62
Stroke or TIA	6 (1%)	4 (1%)	0.69 (0.19-2.44)	0.57
Urgent revascularisation	8 (1%)	9 (1%)	1.17 (0.45-3.04)	0.74
Safety endpoints				
STEEPLE major bleed	7 (1%)	1 (<0.5%)	0.15 (0.02-1.20)	0.07
STEEPLE minor bleed	5 (1%)	2 (<0.5%)	0.41 (0.08-2.13)	0.29
STEEPLE major or minor bleed	12 (2%)	3 (1%)	0.26 (0.07-0.91)	0.04

Data are n (%) unless otherwise stated. *All three stent thromboses were definite. TIA=transient ischaemic attack. TIMI=thrombolysis in myocardial infarction.

Table 3: Study endpoints during follow-up

Outcomes

The main study objective was to show the superiority of continuation of DAPT beyond 1 year over a strategy of stopping thienopyridine. The primary endpoint was a composite of all-cause death, myocardial infarction, stroke or transient ischaemic attack, urgent coronary revascularisation, and stent thrombosis, the same endpoint as for the first phase of the study analysed by intention to treat. All definitions are described elsewhere.^{21,22} The main secondary efficacy endpoint was the composite of stent

thrombosis (whether revascularised or not) and urgent revascularisation. We also analysed all the other protocol-specified study endpoints. The main safety endpoint was defined as major bleeding using the percutaneous

coronary intervention-specific STEEPLE definitions.²⁴ All events were adjudicated by an independent clinical events committee (Pitié-Salpêtrière University Hospital, Institut de Cardiologie, Paris, France) unaware of treatment assignments. We assessed all interactions between high platelet reactivity during treatment with thienopyridine, defined as 235 or more platelet reaction units and treatment group for all pre-specified endpoints.

Statistical analysis

We hypothesised that the ischaemic event rate would be 6% within the first 6 months of interruption of DAPT. We calculated that we would need a sample size of 1492 patients to allow an 80% power to detect a 50% relative risk reduction by a 5% two-sided survival test based on a Cox model. The analysis was based on all events that occurred in the intention-to-treat population, defined as all randomly allocated patients who signed an informed consent form. In case of patients withdrawing consent during the study, only their data collected before the day of withdrawal were included in the database. We analysed the endpoints using a Cox model for survival analysis. We also tested outcomes for their association with high on-treatment platelet reactivity by linear regression using a codominant model after adjusting for clinical characteristics associated with high platelet reactivity at an α -level of <0.05 . All patients were censored at the date of last available information. We summarised non-Gaussian variables as median (IQR) and compared them with the Mann-Whitney test. We used the χ^2 test for frequency comparisons. All subgroups presented were pre-specified. All tests were made at a two-sided 5% significance level. We also did on-treatment analyses by taking treatment exposition as a time-dependent covariate to assess the effect of patients who crossed over. We used SAS (version 9.2) for all statistical analyses.

The study protocol was registered at ClinicalTrials.gov, number NCT00827411.

Role of the funding source

The trial and the statistical analyses were done by the non-for-profit academic research organisation ACTION (Allies in Cardiovascular Trials, Initiatives and Organized Networks), based at Pitié-Salpêtrière Hospital). Research grants were obtained from Fondation de France, Sanofi-Aventis, Cordis, Medtronic, Boston Scientific, Fondation SGAM which had no involvement in the conduct of the study. GM and J-PC had full access to data and full responsibility for the decision to submit for publication.

Results

Of the 2440 patients originally enrolled in the ARCTIC-TRIAL, 1259 were enrolled into the ARCTIC-Interruption study, of whom 624 were assigned to the interruption group and 635 were assigned to the continuation group (figure 1). The baseline risk of the non-randomised study population (n=1181) differed




Figure 2: Kaplan-Meier curves for the primary (A) and main secondary (B) endpoints and main safety endpoints (C)
 (A) The primary endpoint was death, myocardial infarction, stent thrombosis, stroke, or urgent revascularisation.
 (B) The main secondary endpoint was stent thrombosis or any urgent revascularisation.
 (C) The main safety endpoint was major and minor STEEPLE bleedings.




Figure 3: Occurrence of primary (A) and secondary (B) endpoints in prespecified subgroups
ACS=acute coronary stenting. BMI=body-mass index.

from the ARCTIC-Interruption population with a higher proportion of diabetes (469 [40%] of 1181 patients *vs* 420 [33%] of 1259 patients; $p=0.0011$), peripheral artery disease (167 [14%] patients *vs* 124 [10%] patients; $p=0.0011$; appendix), and higher platelet reactivity to adenosine diphosphate measured on thienopyridine maintenance treatment before interruption (158 [SD 92] *vs* 145 [88]; $p=0.014$). The main reasons for non-randomisation were physician's decision ($n=671$ [57%]), time from first randomisation greater than 15 months ($n=253$ [21%]), occurrence of major cardiovascular events or an absolute indication of extended DAPT (eg, new revascularisation; $n=201$ [17%]), lack of treatment adherence ($n=29$ [2%]), and the need for chronic oral anticoagulation ($n=27$ [2%]), baseline characteristics are reported in table 1). Use of antiplatelet treatment at the time of re-randomisation was well balanced between study groups except for clopidogrel 150 mg (table 2). Roughly 10% of patients were on prasugrel maintenance treatment and roughly 10% were on 150 mg clopidogrel maintenance treatment, with both treatments necessary because of previous treatment adjustment in patients initially randomly allocated to platelet function testing in the first phase of the study.

During the follow-up period, adherence to the assigned DAPT treatment progressively decreased in the continuation group while in the interruption group some patients had to be restarted on DAPT (table 2). Nevertheless, there was still a large difference between the two groups (table 2).

During the follow-up period of 17 months (IQR 15–18), 16 (1%) patients died, 18 (1%) had an acute myocardial infarction, 17 (1%) had an urgent revascularisation, three (1%) had a definite or probable stent thrombosis, and ten (1%) had a stroke. At the end of follow-up, we saw no between-group difference in the primary endpoint (table 3, figure 2). This finding was consistent for all the secondary endpoints (table 3, figure 2). We also saw no between-group differences across any subgroups for both the primary and main secondary endpoints (figure 3). High platelet reactivity to ADP during thienopyridine treatment prior to randomisation was associated with a higher mortality rate at the end of follow-up than that in patients without high platelet reactivity (six [3%] of 181 patients *vs* one [1%] of 181 patients; hazard ratio 5.07 [95% CI 1.63–15.76]; $p=0.005$). However, we detected no interaction between platelet reactivity to ADP and interruption or continuation of

See Online for appendix

For more on ACTION
see <http://www.action-coeur.org>

thienopyridine for the primary study endpoint ($p=0.78$; appendix) or for any other secondary ischaemic endpoint.

Overall, there were 15 bleeding events according to the STEEPLE definitions, of which eight were major and seven were minor (table 3). There was no statistically significant between-group difference in major bleeds (hazard ratio with interruption 0.15 [95% CI 0.02–1.20]; table 3). Analysis of both major and minor bleeds combined showed bleeding to be less frequent in the interruption group (0.26 [95% CI 0.07–0.91]; figure 2, table 3). No patient had a TIMI major bleed or a fatal bleed. We saw no interaction between platelet reactivity to ADP measured before randomisation and treatment group for safety outcomes (appendix). Sensitivity analyses taking into account the timing and the number of crossover patients showed similar results as the intention-to-treat analysis (appendix).

Discussion

Findings from ARCTIC-Interruption show no benefit with DAPT continuation beyond 1 year after coronary stenting, rather a harm of continued DAPT, with

patients in the continuations group having more major or minor bleeding. They also show that 1 year after stenting, only half of patients are eligible for randomisation to thienopyridine interruption or continuation leading to the selection of a low risk population, and that strict adherence to the open-label treatment assignment of both single and dual antiplatelet treatment beyond 1 year after stenting was not possible in a quarter of patients, a rate that is consistent with observations in clinical practice.²⁵

Eight characteristics of the study suggest that our findings are applicable to the general population of patients. First, before being regarded as eligible for interruption, patients underwent a first screening for the ARCTIC study and needed to be in stable condition and a candidate for coronary stenting with a DES.²² Second, the patients were exposed to DAPT for 1 year after DES implantation, a duration recommended by clinical guidelines.^{6,26} Third, only half the ARCTIC population was eligible for the second phase assessing thienopyridine interruption, a finding that is consistent with results of other studies—eg, in the prospective observational PARIS registry,²⁵ physicians discontinued DAPT 1 year after stenting in only 41% of patients. Fourth, by contrast with previous studies,^{18–20} we randomly allocated patients for DAPT interruption not at the time of stenting but 1 year later, which better follows clinical practice. Subsequently, we restricted enrolment to patients who remained event-free 1 year after the index percutaneous coronary intervention and higher risk patients were likely to be excluded. Fifth, the global follow-up of our patients could be extended up to 30 months, a duration longer than what is usually seen in interventional studies. Sixth, event rates in ARCTIC-Interruption were within the ranges expected from registries, outlining that the study is relevant to real-world practice.²⁵ Seventh, the results of the ARCTIC-Interruption study were consistent irrespective of the timing and the number of crossover patients. Finally, the absence of benefit of extended DAPT reported in ARCTIC-Interruption lends support to findings from the randomised CHARISMA trial which was done in stable, at-risk patients who were medically managed (panel).²⁷

Measurement of platelet reactivity was another unique aspect of the ARCTIC-Interruption trial. Although platelet reactivity was shown to be a risk marker in the first phase of the ARCTIC study, antiplatelet treatment adjustment in patients with persistent high on-treatment platelet reactivity did not improve outcome.²² Whether platelet reactivity should be measured in specific situations is under debate.^{30–32} In the ARCTIC-Interruption study, platelet reactivity to ADP was measured on P2Y₁₂ inhibitor maintenance dose and was regarded as a baseline characteristic. The finding that non-randomised patients had higher platelet reactivity substantiates the fact that platelet reactivity is a marker of the global vascular risk of patients. That platelet reactivity did not differ between the continuation and the interruption

Panel: Research in context

Systematic review

To further assess the effect of treatment duration on clinical outcome, we searched PubMed and the Cochrane databases from Jan 1, 2002, to Sept 1, 2013, using the following search terms: “dual antiplatelet therapy”, “aspirin”, “clopidogrel”, “stent(s)”, “drug-eluting stent(s)”, “PCI” (percutaneous coronary intervention), and “randomized trial”. We did not include studies with P2Y₁₂ inhibitors other than thienopyridine and included studies with a minimum follow-up of 3 months from randomisation. On the basis of this review we identified no clear effect of the duration of interruption. Indeed, there was one trial¹⁶ with a 1 year minimum duration of DAPT. We did a meta-analysis of all randomised studies that assessed interruption of dual antiplatelet treatment (DAPT) after coronary stenting with drug-eluting stents (DESs). We included six randomised studies including the ARCTIC-Interruption study with a total of 12 536 enrolled patients, of whom 6252 were assigned to short DAPT duration defined as a minimum of 3 months.^{16–20} Events were independently adjudicated in all studies and randomisation took place at the time of index PCI,^{18–20} or 1 month¹⁷ or 1 year later.¹⁶ All cause death and major bleeding were the primary objectives of this meta-analysis. The endpoint definitions were those of each individual study. Two investigators (JPC and JS) independently assessed reports for eligibility on review of the title or the abstract (appendix). The study was done in compliance with the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) statement (appendix) and the protocol is available on request.

Interpretation

When pooling all the evidence, at a median follow-up of 17 months, continuation of DAPT did not reduce reduced all-cause mortality (figure 4), major adverse cardiac events (1.05 [95% CI 0.87–1.25]; $p=0.62$), or the net clinical benefit (1.07 [0.89–1.27]; $p=0.48$). None of the individual ischaemic endpoints was reduced by continuation of DAPT, including stent thrombosis (0.86 [0.53–1.39]; $p=0.41$), stroke (1.43 [0.93–2.21]; $p=0.10$), and myocardial infarction (1.03 [0.79–1.34]; $p=0.84$). Major bleeding was increased by two times (figure 4). These findings were consistent irrespective of the DAPT duration in the interruption group. Additionally, we saw no statistically significant heterogeneity and the sample size of this meta-analysis is in the range of the large ongoing trials assessing the effect of DAPT interruption (NCT00661206, NCT00977938).^{28,29}




Figure 4: Outcomes according to continuation or interruption of dual antiplatelet treatment after percutaneous coronary interventions

Data are for death (A) or major bleeding (B). Analyses were dichotomised according to the minimum duration of dual antiplatelet treatment in the control group. M-H Random=Mantel-Haenszel Random. p_{het}=p for heterogeneity.

groups suggests that not only antiplatelet treatment but also response to treatment were well balanced in the two groups after randomisation in ARCTIC-Interruption. In this randomised population, we saw a strong relation between baseline platelet reactivity to ADP and survival, substantiating once more the prognostic value of this marker. However, the absence of statistically significant interaction between interruption and platelet reactivity on clinical outcomes validates further the main results of the first phase of the ARCTIC study. The results of ARCTIC-Monitoring and ARCTIC-Interruption emphasise that platelet reactivity is of restricted value to guide decisions of augmentation or interruption of antiplatelet treatment.

We acknowledge that the ARCTIC-Interruption trial enrolled a sample size smaller than anticipated, used power assumptions based on data from the era of first-generation of DESs, and did not enrol patients at very high thrombotic burden. These limitations further draw attention to the need for confirmatory data from ongoing larger studies with longer follow-up, stronger

pharmacological intervention, and higher-risk patients (NCT00977938, NCT01225562).^{33,34} Additionally, the absence of interaction between interruption and platelet reactivity on clinical outcomes should be viewed as merely hypothesis-generating.

The randomised ARCTIC-Interruption study and the meta-analysis suggest no apparent benefit, but instead harm, with extension of DAPT beyond 1 year after stenting with DES when no event has occurred within the first year of treatment.

Contributors

GM, J-PC, and EV designed the study and analysed and interpreted the data. GC, TC, SE, GR, J-PC, GM participated to the steering committee, contributed to implementation of the study, enrolment, and follow-up of patients, and revised the manuscript. HR and EV did all statistical analyses. GM and J-PC wrote the first draft and submitted the final version for publication. TC, GR, GC, SE, PH, DC, LB, EV, PA, JM, PS, SAO'C, JA, MK, CS-E, OB, FB, and JS participated to the study, commented on the manuscript, and approved the final version. All authors have seen the version of the Article and agree with the content and conclusions.

Declaration of interests

J-PC reports receiving research grants from Bristol-Myers Squibb, Sanofi-Aventis, Eli Lilly, Medtronic, Boston Scientific, Cordis, Stago, Fondation de France, INSERM, Nanospheres, Fédération Française de Cardiologie and Société Française de Cardiologie; consulting fees from Sanofi-Aventis, Eli Lilly, and Bristol-Myers Squibb; and lecture fees from Bristol-Myers Squibb, Sanofi-Aventis, Eli Lilly, and AstraZeneca. TC reports receiving a research grant from Eli Lilly; consulting fees from Daiichi Sankyo and Eli Lilly; support for travel to meetings from Eli Lilly Daiichi Sankyo; lecture fees from AstraZeneca, Eli Lilly, Sanofi, Abbott Vascular, Biotronik, Boston Scientific, Cordis, Edwards, Iroko Cardio, Medtronic, Servier; travel and meetings expenses from Abbott Vascular, AstraZeneca, Biotronik, Boston Scientific, Cordis, Daiichi Sankyo, Edwards, Eli Lilly, Iroko Cardio, Medtronic, and Servier. GC reports receiving a research grant from la Fédération Française de Cardiologie; consultant fees from Abbott Vascular, AstraZeneca, CLS, Behring, Daiichi Sankyo, and Eli Lilly; and lecture fees from Abbott Vascular, AstraZeneca, Biotronik, CLS Behring, Daiichi Sankyo, Eli Lilly, and Iroko Cardio. CP reports receiving research support and consulting fees from Sanofi-Aventis and The Medicines Company. PM has received consultancy fees from St Jude Medical and Terumo. PS reports receiving fees for board membership from Bristol-Myers Squibb; consulting fees from Daiichi Sankyo and Eli Lilly; and lecture fees from Merck Sharp and Dohme and Sanofi. SAO'C reports receiving grants from A Menarini and the European Society of Cardiology. JA reports receiving a grant from Fonds d'Etudes et de Recherche du Corps Médical (FERCM). MK reports receiving a grant from Fédération Française de Cardiologie. FB reports receiving grants from Pfizer, Brahms, and Fédération Française de Cardiologie; and lecture fees from Roche, Sanofi-Aventis, Pfizer, and Astellas. JS reports receiving research grants from Sanofi-Aventis, Daiichi-Sankyo, Eli Lilly, Brahms, INSERM, Fédération Française de Cardiologie, and Société Française de Cardiologie; consulting fees from Daiichi-Sankyo and Eli Lilly; and speakers' honoraria from AstraZeneca, Daiichi Sankyo, Eli Lilly, and Servier. EV reports receiving research grants from Sanofi-Aventis, Medtronic, Cordis, Boston Scientific, Fondation SGAM, Fondation de France, ACTION and APHP; consulting fees from Pfizer, Eli Lilly, Sanofi, LFB, Abbott, Fresenius; grant from Boehringer-Ingelheim; and lecture fees from Merck and Pfizer. GM reports receiving grant support from Abbott Vascular, Boston Scientific, Cordis, Eli Lilly, Fédération Française de Cardiologie, Fondation de France, Guerbet Medical, INSERM, ITC Edison, Medtronic, Pfizer, Sanofi-Aventis, Société Française de Cardiologie, and Stago; consulting or board fees and lecture fees from AstraZeneca, Bayer, Boehringer-Ingelheim, Cardiovascular Research Foundation, Cleveland Clinic Research Foundation, Daiichi-Sankyo, Duke Institute, Eli Lilly, Europa, Lead-up, GlaxoSmithKline, Institut de Cardiologie de Montréal, Menarini, Nanospheres, Novartis, Pfizer, Portola, Sanofi-Aventis, The Medicines Company, and the TIMI study group.

Acknowledgments

We thank Yannick Vacher, Abdel Ait Bachir, Andrea Bardon, Nabil Brouk, Julie Bussone, Hermione Folal, Vanessa Gallois, Sophie Gérard, Véronique Jouis, Azizath Kabirou, Nathalie Kingue, Valérie Mazur, Hélène Mauro, Ghislain N'Gouala, Luminita Neculaita, Sissel Paulsrud, Virginie Rochaud, and Corinne Tchokothe.

References

- Task Force Members, Montalescot G, Sechtem U, et al. 2013 ESC guidelines on the management of stable coronary artery disease: The Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J* 2013; **34**: 2949–3003.
- Daemen J, Wenaweser P, Tsuchida K, et al. Early and late coronary stent thrombosis of sirolimus-eluting and paclitaxel-eluting stents in routine clinical practice: data from a large two-institutional cohort study. *Lancet* 2007; **369**: 667–78.
- Eisenstein EL, Anstrom KJ, Kong DF, et al. Clopidogrel use and long-term clinical outcomes after drug-eluting stent implantation. *JAMA J Am Med Assoc* 2007; **297**: 159–68.
- Pfisterer M, Brunner-La Rocca HP, Buser PT, et al. Late clinical events after clopidogrel discontinuation may limit the benefit of drug-eluting stents: an observational study of drug-eluting versus bare-metal stents. *J Am Coll Cardiol* 2006; **48**: 2584–91.
- Morice M-C, Serruys PW, Sousa JE, et al. A randomized comparison of a sirolimus-eluting stent with a standard stent for coronary revascularization. *N Engl J Med* 2002; **346**: 1773–80.
- Wijns W, Kolh P, Danchin N, et al. Guidelines on myocardial revascularization: The Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J* 2010; **31**: 2501–55.
- Kastrati A, Mehilli J, Pache J, et al. Analysis of 14 trials comparing sirolimus-eluting stents with bare-metal stents. *N Engl J Med* 2007; **356**: 1030–39.
- Baber U, Mehran R, Sharma SK, et al. Impact of the everolimus-eluting stent on stent thrombosis: a meta-analysis of 13 randomized trials. *J Am Coll Cardiol* 2011; **58**: 1569–77.
- Sarno G, Lagerqvist B, Fröbert O, et al. Lower risk of stent thrombosis and restenosis with unrestricted use of 'new-generation' drug-eluting stents: a report from the nationwide Swedish Coronary Angiography and Angioplasty Registry (SCAAR). *Eur Heart J* 2012; **33**: 606–13.
- Sabate M, Cequier A, Iñiguez A, et al. Everolimus-eluting stent versus bare-metal stent in ST-segment elevation myocardial infarction (EXAMINATION): 1 year results of a randomised controlled trial. *Lancet* 2012; **380**: 1482–90.
- Airoldi F, Colombo A, Morici N, et al. Incidence and predictors of drug-eluting stent thrombosis during and after discontinuation of thienopyridine treatment. *Circulation* 2007; **116**: 745–54.
- Schulz S, Schuster T, Mehilli J, et al. Stent thrombosis after drug-eluting stent implantation: incidence, timing, and relation to discontinuation of clopidogrel therapy over a 4-year period. *Eur Heart J* 2009; **30**: 2714–21.
- Kandzari DE, Barker CS, Leon MB, et al. Dual antiplatelet therapy duration and clinical outcomes following treatment with zotarolimus-eluting stents. *JACC Cardiovasc Interv* 2011; **4**: 1119–28.
- Hahn J-Y, Song YB, Choi J-H, et al. Three-month dual antiplatelet therapy after implantation of zotarolimus-eluting stents: the DATE (Duration of Dual Antiplatelet Therapy After Implantation of Endeavor Stent) registry. *Circ J Off J Jpn Circ Soc* 2010; **74**: 2314–21.
- Naidu SS, Krucoff MW, Rutledge DR, et al. Contemporary incidence and predictors of stent thrombosis and other major adverse cardiac events in the year after XIENCE V implantation: results from the 8,061-patient XIENCE V United States study. *JACC Cardiovasc Interv* 2012; **5**: 626–35.
- Park S-J, Park D-W, Kim Y-H, et al. Duration of dual antiplatelet therapy after implantation of drug-eluting stents. *N Engl J Med* 2010; **362**: 1374–82.
- Valgimigli M, Campo G, Monti M, et al. Short- versus long-term duration of dual-antiplatelet therapy after coronary stenting: a randomized multicenter trial. *Circulation* 2012; **125**: 2015–26.
- Kim B-K, Hong M-K, Shin D-H, et al. A new strategy for discontinuation of dual antiplatelet therapy: the RESET Trial (REal Safety and Efficacy of 3-month dual antiplatelet Therapy following Endeavorzotarolimus-eluting stent implantation). *J Am Coll Cardiol* 2012; **60**: 1340–48.
- Gwon H-C, Hahn J-Y, Park KW, et al. Six-month versus 12-month dual antiplatelet therapy after implantation of drug-eluting stents: the Efficacy of Xience/Promus Versus Cypher to Reduce Late Loss After Stenting (EXCELLENT) randomized, multicenter study. *Circulation* 2012; **125**: 505–13.
- Feres F, Costa RA, Abizaid A, et al. Three vs twelve months of dual antiplatelet therapy after zotarolimus-eluting stents: the OPTIMIZE randomized trial. *JAMA J Am Med Assoc* 2013; **310**: 2510–22.
- Collet JP, Cayla G, Cuisset T, et al. Randomized comparison of platelet function monitoring to adjust antiplatelet therapy versus standard of care: rationale and design of the assessment with a double randomization of (1) a fixed dose versus a monitoring-guided dose of aspirin and clopidogrel after DES implantation, and (2) treatment interruption versus continuation, 1 year after stenting (ARCTIC) study. *Am Heart J* 2011; **161**: 5–12.
- Collet J-P, Cuisset T, Rangé G, et al. Bedside monitoring to adjust antiplatelet therapy for coronary stenting. *N Engl J Med* 2012; **367**: 2100–09.
- Silber S, Albertsson P, Aviles FF, et al. Guidelines for percutaneous coronary interventions. The task force for percutaneous coronary interventions of the European Society of Cardiology. *Eur Heart J* 2005; **26**: 804–47.

- 24 Montalescot G, White HD, Gallo R, et al. Enoxaparin versus unfractionated heparin in elective percutaneous coronary intervention. *N Engl J Med* 2006; **355**: 1006–17.
- 25 Mehran R, Baber U, Steg PG, et al. Cessation of dual antiplatelet treatment and cardiac events after percutaneous coronary intervention (PARIS): 2 year results from a prospective observational study. *Lancet* 2013; **382**: 1714–22.
- 26 Hamm CW, Bassand J-P, Agewall S, et al. ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: The Task Force for the management of acute coronary syndromes (ACS) in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2011; **32**: 2999–3054.
- 27 Bhatt DL, Fox KA, Hacke W, et al. Clopidogrel and aspirin versus aspirin alone for the prevention of atherothrombotic events. *N Engl J Med* 2006; **354**: 1706–17.
- 28 Byrne RA, Schulz S, Mehilli J, et al. Intracoronary stenting and antithrombotic regimen: safety and efficacy of six months dual antiplatelet therapy after drug-eluting stenting (ISAR-safe) investigators rationale and design of a randomized, double-blind, placebo-controlled trial of 6 versus 12 months clopidogrel therapy after implantation of a drug-eluting stent: the intracoronary stenting and antithrombotic regimen: safety and efficacy of 6 months dual antiplatelet therapy after drug-eluting stenting (ISAR-SAFE) study. *Am Heart J* 2009; **157**: 620–24.
- 29 Mauri L, Kereiakes DJ, Normand S-LT, et al. Rationale and design of the dual antiplatelet therapy study, a prospective, multicenter, randomized, double-blind trial to assess the effectiveness and safety of 12 versus 30 months of dual antiplatelet therapy in subjects undergoing percutaneous coronary intervention with either drug-eluting stent or bare metal stent placement for the treatment of coronary artery lesions. *Am Heart J* 2010; **160**: 1035–41.
- 30 Aradi D, Storey RF, Komocsi A, et al. Expert position paper on the role of platelet function testing in patients undergoing percutaneous coronary intervention. *Eur Heart J* 2013; **35**: 209–15.
- 31 Tantry US, Bonello L, Aradi D, et al. Consensus and update on the definition of on-treatment platelet reactivity to adenosine diphosphate associated with ischemia and bleeding. *J Am Coll Cardiol* 2013; **62**: 2261–73.
- 32 Montalescot G, Vicaut E, Collet J-P. Bedside monitoring of antiplatelet therapy for coronary stenting. *N Engl J Med* 2013; **368**: 871–72.
- 33 Bonaca MP, Bhatt DL, Braunwald E, et al. Design and rationale for the prevention of cardiovascular events in patients with prior heart attack using ticagrelor compared to placebo on a background of aspirin-thrombolysis in myocardial infarction 54 (PEGASUS-TIMI 54) trial. *Am Heart J* 2014; **167**: 437–44.
- 34 Mauri L, Kereiakes DJ, Normand S-LT, et al. Rationale and design of the dual antiplatelet therapy study, a prospective, multicenter, randomized, double-blind trial to assess the effectiveness and safety of 12 versus 30 months of dual antiplatelet therapy in subjects undergoing percutaneous coronary intervention with either drug-eluting stent or bare metal stent placement for the treatment of coronary artery lesions. *Am Heart J* 2010; **160**: 1035–41.